

Drug Information Sheet("Kusuri-no-Shiori")

Injection

Published: 04/2018

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:ROCEPHIN I.V. 1g

Active ingredient:Ceftriaxone sodium hydrate

Dosage form:injection

Print on wrapping:



Effects of this medicine

This medicine kills bacteria which cause infectious disease or inflammation.

It is usually used for the treatment of sepsis, pharyngitis/laryngitis, tonsillitis, acute bronchitis, pneumonia, lung abscess, pyothorax, secondary infection of chronic respiratory lesion, cystitis, pyelonephritis, epididymo-orchitis (epididymitis), urethritis, cervicitis, pelvic inflammatory disease, proctitis, peritonitis, intraperitoneal abscess, cholecystitis, cholangitis, bartholinitis, intrauterine infection, adnexitis, parametritis, purulent meningitis, keratitis (including corneal ulcer), otitis media, sinusitis, perimaxillary cellulitis and jaw inflammation.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
If you or any of your parents or siblings predisposed to allergic reactions, such as bronchial asthma, rash, and urticaria.
- If you have renal disorder, poor food intake or physical debility.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, this medicine is given once daily or twice daily for adults and children, once daily for premature infants and neonates aged 0-3 days, twice daily for premature infants and neonates aged 4 days or older, by intravenous injection. It can be used by intravenous drip infusion.
- The treatment period with this medicine is decided, depending on the symptoms.

Precautions while taking this medicine

- The medicine may cause deficiency of vitamin B, K, etc., which may result in symptoms such as nose bleeding tendency or stomatitis. Ingest enough foods to prevent vitamin deficiency. Contact your doctor when you cannot ingest enough foods because of loss of appetite, etc.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include rash, diarrhea, hives, fever, redness, itch and erythema. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- cold sweat, dizziness on standing up, breathing difficulty [shock, anaphylactoid symptoms]
- general malaise, easy bruising, easy to bleed (gum/nose bleeding, prolonged menstruation) [pancytopenia, agranulocytosis, white blood cell decreased, platelets decreased, haemolytic anaemia]
- fever, decreased urinary volume, edema [renal failure acute, nephritis interstitial]
- abdominal pain, frequent diarrhea [pseudomembranous colitis]
- mucosal swelling in the eyes/lips/genitalia, widespread redness in the skin, blister [toxic epidermal necrolysis, oculomuocutaneous syndrome, acute generalized eruptive pustulosis]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.